

BOTANISKA NOTISER

FÖR ÅR 1962

VOL. 115

UTGIVEN AV

LUNDS BOTANISKA FÖRENING

REDIGERAD AV

H. HJELMQVIST

Tidskriften grundad 1839

DISTRIBUTÖR: C. W. K. GLEERUP, FÖRLAG · LUND

Utgivningstider:

- Fasc. 1, sid. 1—112, den 29 mars
Fasc. 2, sid. 113—244, den 28 juni
Fasc. 3, sid. 245—356, den 29 sept.
Fasc. 4, sid. 357—513, den 29 dec.

LUND 1962

CARL BLOMS BOKTRYCKERI A.-B.

INNEHÅLLSFÖRTECKNING

ALMQUIST, ERIK: Ett par skånska hieracier av dansk härkomst	101
ARNELL, SIGFRID: Hepaticae collected by Dr. O. H. Selling in Central Australia, Tasmania and New Zealand in 1949	311
— Notes on South African Hepaticae VII	203
BERGLUND, BJÖRN, Vegetation på ön Senoren. I. Vegetationshistoria. (Summary p. 416: Vegetation of the island of Senoren. I. Vegetational history.)	387
BOULOS, LOUTFY: Cytotaxonomic studies in the genus <i>Sonchus</i> . 4. The generic status of some species earlier treated as <i>Sonchus</i>	58
BRINGER, KARL-GÖRAN: Kärleväxtfloran i en serie sydexponerade branter och brantavsnitt på fjället Vakkettjåkko, Torne lappmark, samt några iakttagelser rörande sydväxtbergfloras ståndortsekologi. (Summary p. 339: The vascular plants of south facing bluffs on Vakkettjåkko, a mountain in the Torneträsk area, northern Swedish Lappland, and some observations concerning their distribution and ecology.)	318
CHRTEK, JINDŘICH, and KRÍSA, BOHDAN: A taxonomical study of the species <i>Luzula spicata</i> (L.) DC. sensu lato in Europe	293
DAHLGREN, K. V. OSSIAN: Die Lösung des micrantha-Problems bei <i>Fragaria vesca</i> . (Summary p. 291: The solving of the micrantha-problem of <i>Fragaria vesca</i> .)	288
DAHLGREN, ROLF: Some new species of <i>Aspalathus</i> from the Cape Province	465
DIXON, PETER S.: Taxonomic and nomenclatural notes on the Florideae, III	245
ENGLESSON, NILS: Nya lokaler för <i>Senecio erucifolius</i> och <i>Dianthus superbus</i> ..	477
FLINCK, KARL EVERT, and HYLMÖ, BERTIL: <i>Cotoneaster Harrysmithii</i> , a new species from Western China	29
— <i>Cotoneaster sikangensis</i> , a new species from Western China	376
— — On two recently described species of <i>Cotoneaster</i> of northwestern Europe	343
GOLDSCHMIDT, ERICH, se NILSSON, FREDRIK.	
HALLECK, DIANNE, se WIENS, DEL.	
HANSEN, ALFRED: <i>Dipsacus pilosus</i> L. findes næppe i Sverige. (Zusammenfassung S. 107: <i>Dipsacus pilosus</i> ist kaum in Schweden vorhanden.)	106
HARRIS, R. E.: Contribution to the taxonomy of <i>Callithamnion Lyngbye</i> emend. Naegeli	18
HJELMQVIST, H.: A case of virescence in <i>Trifolium hybridum</i>	429
— The embryo sac development of some <i>Cotoneaster</i> species	208
HOLMEN, KJELD: Chromosome studies in some Arctic Alaskan Leguminosae	87
HYLMÖ, BERTIL, se FLINCK, KARL EVERT.	
JEFFREY, C.: The origin and differentiation of the archegoniate land-plants	446
KRÍSA, BOHDAN, se CHRTEK, JINDŘICH.	
LARSEN, KAI: Contribution to the cytology of the endemic Canarian element	196
LÖVE, ÅSKELL: Typification of <i>Papaver radicatum</i> — a nomenclatural detective story	113
LÖVKVIST, BÖRJE: A case of probable introgression in <i>Cirsium</i>	385
— Chromosome and differentiation studies in flowering plants of Skåne, South Sweden. I. General aspects. Type species with coastal differentiation	261

NILSSON, ARVID: <i>Juncus maritimus</i> vid Skanör. En ny medborgare i Skånes flora	102
NILSSON, FREDRIK, and GOLDSCHMIDT, ERICH: Cytochimeras in <i>Ribes</i>	137
NILSSON, SVEN: Second note on Swedish freshwater Hyphomycetes	73
NORLINDH, TYCHO: Studies in <i>Calendula maderensis</i> DC. With a discussion on the delimitation of <i>Calendula</i> L. from <i>Gibbaria</i> Cass. and <i>Osteospermum</i> L.	437
PETTERSSON, BENGT: Om <i>Ceterach officinarum</i> och dess ekologi	237
RECHINGER, K. H.: Revision einiger Typen von Velenovsky's <i>Plantae Arabicae Musiliana</i> e	35
RICKMAN, HELGE: Några skånska växtnotiser	475
RUNEMARK, HANS: A revision of <i>Parapholis</i> and <i>Monerma</i> in the Mediterranean	1
— A revision of <i>Pteris dentata</i> and related species	177
RUNGBY, SVEND: A contribution to the bryophytic flora of Spain and Morocco, especially the area between Gandia and Alcoy	61
SAAD, SHOKRY I.: Pollen morphology of Ctenolophon	49
SAARSOO, BERNHARD: Tre nya <i>Taraxacum</i> -arter från Finland. (Summary p. 72: Three new species of <i>Taraxacum</i> from Finland.)	65
SAHLIN, C. I.: <i>Dryopteris abbreviata</i> (DC.) Newm. från Lofoten	240
— <i>Vaccinium oxycoccus</i> L. (s. str.), ny för Torne lappmark	480
SNOGERUP, SVEN: Studies in the Aegean flora. IV. <i>Bupleurum flavum</i> Forsk. and related species	357
TRALAU, HANS: Die spättertiären <i>Fagus</i> -Arten Europas. (Summary p. 169: The Late-Tertiary <i>Fagus</i> species of Europe.)	147
— <i>Najas tenuissima</i> (A. Br.) Magnus during the Late-Cainozoic period in Europe	421
WIENS, DEL, and HALLECK, DIANNE K.: Chromosome numbers in Rocky Mountain plants. I	455

Lunds Botaniska Förening

Från Lunds Botaniska Förenings förhandlingar 1961	93
Lunds Botaniska Förening 1962	511

Litteratur

Svensk Botanisk Litteratur 1961. (Av ARNE H. HOLMQVIST.)	481
Atlas of the British Flora. Ed. by F. H. PERRING and S. M. WALTERS. (Av H. HJELMQVIST.)	354
CLAPHAM, A. R., TUTIN, T. G., and WARBURG, E. F.: Flora of the British Isles. 2nd ed. (Av H. HJELMQVIST.)	354
EAMES, ARTHUR J.: Morphology of the angiosperms. (Av H. HJELMQVIST.) ..	507
GUPTA, K. M.: Marsilea. (Av H. HJELMQVIST.)	508
HAWKER, LILIAN E., LINTON, A. H., FOLKES, B. F., and CARLILE, M. J.: An introduction to the biology of micro-organisms. (Av B. NORÉN.)	109
HUBER-PESTALOZZI, GOTTFRIED: Das Phytoplankton des Süßwassers. Teil 5. Chlorophyceae (Grünalgen). Ordn. Volvocales. (Av EINAR TEILING.)	352
MAHESHWARI, P., and VASIL, VIMLA: Gnetum. (Av H. HJELMQVIST.)	508
MALMGREN, BERNDT: Introduktion till mikrobiologien. (Av B. NORÉN)	109
Nye tidsskrifter. Phytochemistry. Photochemistry. Radiation Botany. Life Sciences. (Av PER HALLDAL.)	351
VASILJEV, I. M.: Wintering of plants. (Av BENGT LÖÖF.)	110
Notiser	112, 242, 355, 509

REGISTER

Nedanstående förteckning upptager endast sådana taxa, som blivit i något avseende utförligare eller mera speciellt behandlade. Nya arter och kombinationer betecknas med fetstil. * anger att avbildning, k att karta förekommer.

- | | | | |
|---|-----------|---|------------------|
| <i>Achillea fragrantissima</i> | 46 | <i>Beta patellaris</i> | 197 f* |
| <i>Aconitum columbianum</i> | 458* f | <i>Bonnemaisonia clavata</i> | 256 f |
| <i>Acromastigium integrifolium</i> | 313 | <i>Bupleurum</i> aira | 357 ff* |
| <i>Actinospora megalospora</i> | 73 f* | — <i>flavum</i> | 357 ff* k |
| <i>Aglaothamnion</i> | 19 ff | — <i>glumaceum</i> | 357 ff* k |
| <i>Allagopappus dicotomus</i> | 198*, 201 | — <i>gracile</i> | 357 ff* k |
| <i>Allium cernuum</i> | 457* f | <i>Calendula arvensis</i> | 438 ff* |
| — <i>geyeri</i> | 457* f | — <i>maderensis</i> | 437 ff* |
| <i>Alyssum homalocarpum</i> | 37 | — <i>maritima</i> | 438 ff |
| — (<i>Odontarrhena</i>) <i>singarense</i> | 37 | <i>Callithamnion</i> | 18 ff |
| <i>Anguillospora crassa</i> | 74 f* | <i>Caltha leptosepala</i> | 458* f |
| <i>Anisosciadium lanatum</i> | 42 | <i>Canarina campanula</i> | 198*, 200 |
| — <i>orientale</i> | 42 f | <i>Cephaloziella</i> triplicata | 203 |
| <i>Anthyllis vulneraria</i> ssp. <i>lapponica</i> | 330 f | <i>Ceramium ciliatum</i> | 252 f |
| <i>Arctostaphylos uva-ursi</i> | 457 f* | — <i>pennatum</i> | 254 |
| <i>Arenaria obtusiloba</i> | 456 f* | — <i>robustum</i> | 252 ff |
| <i>Artemisia scopulorum</i> | 457* f | <i>Ceterach officinarum</i> | 237 ff* |
| <i>Articulospora tetracladia</i> | 75* f | <i>Charales</i> | 448 f |
| <i>Aspalathus</i> acanthiloba | 456 f | <i>Chenopodium vulvaria</i> | 476 |
| — campestris | 471 | <i>Chionophila jamesii</i> | 459 f* |
| — linguiloba | 470 f* | <i>Chondria coerulescens</i> | 254 f |
| — lotiflora | 469 f | <i>Cirsium heterophyllum</i> | 385 f |
| — millefolia | 469 | — <i>oleraceum</i> | 385 f |
| — <i>opaca</i> ssp. rostriloba | 473 | <i>Claytonia lanceolata</i> | 458* |
| — pigmentosa | 466 f | <i>Cotoneaster acutifolia</i> v. <i>villosula</i> | 224 ff* |
| — rubiginosa | 467 f* | — <i>Antoninae</i> | 343 ff* |
| <i>Aspicarpa rosei</i> | 52 ff* | — <i>bullata</i> | 216 ff* |
| <i>Astragalus alpinus</i> | 87 f* | — cinerascens | 383 |
| — <i>lepagei</i> | 88 f* | — <i>cinnabarinus</i> | 343 ff* |
| — <i>polaris</i> | 88 f* | — elegans | 383 |
| — <i>umbellatus</i> | 88 f* | — Harrysmithii | 29 ff* |
| — (<i>Ammodendron</i>) <i>camelorum</i> | 39 | — <i>nitens</i> | 31 ff*, 213 ff* |
| — (<i>Platyglottis</i>) <i>bombycinus</i> | 39 | — <i>obscura</i> | 219 ff*, 378 ff* |
| — (<i>Xiphidium</i>) al-hamedensis | 39 f* | — <i>obscura</i> v. <i>nova</i> | 221 ff* |
| Bacillispora aquatica | 77 f* | — <i>racemiflora</i> v. <i>soongorica</i> | 226 ff* |
| <i>Barbula acuta</i> | 61 f | — <i>reticulatus</i> | 379 ff* |
| <i>Bazzania Molleri</i> | 204 | — <i>rosea</i> | 210 ff* |
| — pulvinata | 204 | — sikangensis | 376 ff* |
| <i>Bencomia brachystachya</i> | 198*, 200 | — <i>uniflorus</i> | 343 ff* |
| <i>Besseyia alpina</i> | 459 f* | — villosulus | 383 |

<i>Crouania attenuata</i>	256
<i>Ctenolophon englerianum</i>	49 ff*
— <i>grandifolius</i>	49 ff*
<i>Ctenosiphonia hypnoides</i>	257
<i>Culicidospora</i>	77 f*
<i>Cystopteris fragilis</i> ssp. <i>alpinum</i> ..	332 f
<i>Dactylella microaquatica</i>	78 f*
— submersa	78 f*
<i>Delphinium nelsonii</i>	460* f
<i>Dendrospora erecta</i>	79* f
<i>Dianthus superbus</i>	479
<i>Dicella bracteosa</i>	52 f*
<i>Digitalis isabellina</i>	198*, 201
<i>Dimorphospora foliicola</i>	79* f
<i>Diplotaxis acris</i>	37
<i>Dipsacus pilosus</i>	106 f
— <i>strigosus</i>	106 f
<i>Dryopteris abbreviata</i>	240 f
<i>Echinosciadium arabicum</i>	44
<i>Eremobium nefudicum</i>	38
<i>Erigeron simplex</i>	457* f
<i>Fagus grandifolia</i>	146 ff* k
— <i>silvatica</i>	146 ff* k
<i>Farsetia Burtonae</i>	37
<i>Flagellospora stricta</i>	81* f
<i>Forskahlea angustifolia</i>	198*, 200
<i>Fragaria vesca</i>	288 f*
<i>Galium verum</i>	275 ff*
<i>Gentiana romanzovii</i>	458* f
<i>Geum rossii</i>	460* f
<i>Gibbaria ilicifolia</i>	443 f*
— <i>scabra</i>	443 f*
<i>Gnaphalium luteo-album</i>	475 f
<i>Haplophyllum Blanchei</i>	40 f
<i>Harbouria trachyleura</i>	460* f
<i>Hedysarum alpinum</i>	89* f
<i>Helminthocladia agardhiana</i> ...	245 ff*
<i>Heterocaryum pachypodium</i>	44
<i>Heuchera parvifolia</i>	460* f
<i>Hieracium farumense</i>	101 f
— <i>subastrinum</i>	101 f
— <i>umbellatum</i>	265 ff*
<i>Horwoodia Dicksoniae</i>	36
<i>Hymenoxys acaulis</i>	457* f
— <i>grandiflora</i>	457* f
<i>Hypericum canariense</i>	198*
— <i>grandiflorum</i>	198* f
— <i>reflexum</i>	198* f

<i>Hyalolepidozia bicuspidata</i>	203
<i>Iris spuria</i>	475
<i>Juncus maritimus</i>	102 ff* k
<i>Lastrea oreopteris</i>	477
<i>Launaea Brunneri</i>	59
— <i>Elliotiana</i>	58
— <i>exauriculata</i>	59
— <i>Fischeri</i>	58
— <i>Ledermannii</i>	59
— <i>macer</i>	58
— <i>nanella</i>	58
— <i>pycnocephala</i>	58
— <i>rarifolia</i>	59
— <i>Rueppellii</i>	59
— <i>Verdiekii</i>	58
— <i>violacea</i>	58
<i>Lembidium isodictyon</i>	313
<i>Lemonniera brachycladia</i>	81* f
<i>Lepidozia hippurioides</i>	316
<i>Lewisia pygmaea</i>	458* f
<i>Lloydia serotina</i>	457 f*
<i>Lophocolea excipulata</i>	313
— <i>Sellingii</i>	314*
— <i>turgescens</i>	315
— <i>villosa</i>	315
<i>Lotus corniculatus</i> v. <i>borealis</i> ...	331 f
<i>Lunulospora curvula</i>	83 f*
<i>Lupinus arcticus</i>	91
<i>Luzula bulgarica</i>	304 ff* k
— — ssp. <i>pindica</i>	308 f*
— <i>spicata</i>	293 ff*
— — ssp. <i>spicata</i>	300 ff k
— — ssp. <i>mutabilis</i>	302 ff k
<i>Marchantia pallida</i>	316* f
<i>Maresia pygmaea</i>	37
<i>Matricaria auriculata</i>	45
<i>Matthiola arabica</i>	39
<i>Melanoselinum decipiens</i>	198*, 200
<i>Mertensia lanceolata</i>	457 f*
<i>Mesogloia hudsoni</i>	246 f*
<i>Microlepidozia succulenta</i>	204
— <i>tabularis</i>	204
<i>Monerma cylindrica</i>	2 ff*
<i>Monoclea Forsteri</i>	313
<i>Najas lanceolata</i>	425 f
— <i>tenuissima</i>	421 ff* k
<i>Oreoxis alpina</i> ssp. <i>alpina</i>	460* f
<i>Osteospermum calendulaceum</i> ..	437* f

<i>Oxytropis koyukukensis</i>	88 f*	<i>Riccia Selligii</i>	311 ff*
— <i>leucantha</i>	89* f	<i>Rubus vestitus</i>	477
— <i>maydelliana</i>	89* f	<i>Rumex lunaria</i>	197 f*
— <i>mertensiana</i>	89* f	— <i>maderensis</i>	197 f*
— <i>nigrescens</i>	89* f	<i>Saxifraga adscendens</i>	332
<i>Papaver Dahlianum</i>	116 ff	— <i>flagellaris</i>	459 f*
— <i>Nordhagenianum</i>	128 ff	— <i>rhomboidea</i>	459 f*
— <i>nudicaule</i>	117 ff	<i>Schistochila tuloides</i>	316
— <i>radicatum</i>	113 ff*	<i>Scorzonera Musilii</i>	46
<i>Parapholis filiformis</i>	2 ff*	<i>Sedum rhodanthum</i>	458* f
— <i>incurva</i>	3 ff*	— <i>rosea</i>	456 f*
— marginata	2 ff* k	<i>Senecio congestus</i> v. <i>palustris</i>	476
— <i>pycnantha</i>	7* f k	— <i>erucifolius</i>	477 f
— <i>strigosa</i>	7* f k	— <i>fremontii</i>	457* f
<i>Pedicularis parryi</i>	460* f	— <i>werneriaefolius</i>	458* f
— <i>paysoniana</i>	460* f	<i>Sibbaldia procumbens</i>	459 f*
— <i>racemosa</i>	460* f	<i>Silene acaulis</i>	457*
— <i>scopolorum</i>	460* f	— <i>bourgaei</i>	197 f*
<i>Penstemon virens</i>	460* f	<i>Simia atroviridis</i>	204
<i>Plagiochila lunata</i>	204 f*	<i>Streptopus amplexifolius</i> v. <i>americana</i>	457 f*
— pseudo-attenuata	206*	<i>Taraxacum saetigerum</i>	65 ff*
<i>Plantago amplexicaulis</i> v. <i>bauphula</i>	45	— spurium	68 f*
<i>Polycarpaea carnosa</i> v. <i>carnosa</i>	197 f*	— suberratifrons	68 f*
<i>Potamogeton Friesii</i>	476	<i>Thlaspi arvense</i>	456 f*
<i>Potentilla diversifolia</i>	459 f*	<i>Thymus Musilii</i>	44 f
<i>Prenanthes angustifolia</i>	59	<i>Tortula muralis</i>	62
— suberosa	59	<i>Tradescantia occidentalis</i>	456 f*
<i>Primula angustifolia</i>	458* f	<i>Trichocolea minutifolia</i>	316
— <i>parryi</i>	458* f	<i>Tricladium angulatum</i>	83 f*
<i>Pteris arguta</i>	177 ff* k	— <i>gracile</i>	84 f*
— <i>dentata</i> ssp. <i>dentata</i>	178 ff* k	<i>Trifolium hybridum</i>	429 ff*
— — ssp. flabellata	178 ff* k	— <i>nanum</i>	458* f
— <i>tremula</i>	177 ff*	— <i>parryi</i>	457 f*
<i>Ranunculus adoneus</i>	460* f	<i>Tripleurospermum maritimum</i>	282 ff*
<i>Rectolejeunea richmondensis</i>	315	<i>Vaccinium oxycoccos</i>	480
<i>Rhynia</i>	451	<i>Veronica fruticans</i>	332
<i>Ribes grossularia</i> × <i>divaricatum</i>	138 ff*	<i>Viola rupestris</i> ssp. <i>relicta</i>	330 f
— <i>nigrum</i> × <i>grossularia</i>	138 ff*		

